

CYCADEOIDEA

Kumari Sona Rani
Assistant Professor
Department of Botany
S.Sinha college Aurangabad

For M.sc 2 nd semester

Contents:

1. History of Cycadeoidea
2. Morphological Features of Cycadeoidea
3. Anatomy of Cycadeoidea
4. Reproductive Organs of Cycadeoidea
5. Microsporophyll in Cycadeoidea
6. Gynoecium of Cycadeoidea

1. History of Cycadeoidea:

Cycadeoidea, also called Bennettites by several European palaeobotanists is represented by about 30 species. The name Cycadeoidea was put forward in 1827 for petrified trunks from Isle of Portland. Though Bennettites is still employed for plant fossils from the Isle of Wight, Cycadeoidea is now the valid name of the genus.

It has been reported from Upper Jurassic to Upper Cretaceous rocks of America, India, Russia and several European countries. It occurs in the form of a large number of petrifications in different parts of the world.

2. Morphological Features of Cycadeoidea:

The Cycadeoid trunks were short, stout, spherical to sub-spherical (Figs. 6.1,6.2) and unbranched or branched. The trunks and leaves of many of its species show remarkable resemblance with those of living Cycads. Some of the species were short while others (Cycadeoidea jenneyana) attained a height of 3 to 3.6 metres.

The trunk generally attained a diameter of about 50 cm, and had many, persistent, rhomboidal leaf bases (Figs. 6.2, 6.3). A compact crown of Cycad-like, large, pinnately compound leaves was present at the apex. The leaflets had many parallel veins.




Fig. 6.1. *Cycadeoidea dactyloides*. External features. (after Macbride)




Fig. 6.2. *Cycadeoidea colossalis* showing almost completely spherical stem. (after Wieland).




Fig. 6.3. *Cycadeoidea gibsonianus*. Tangential section through leaf base and flowers.

3. Anatomy of Cycadeoidea:

The stem was roughly circular or oval in outline. It remained covered by heavy armour of leaf bases. The epidermis was not very distinct. The cortex was parenchymatous and possessed many mucilage canals and leaf traces. Many conjoint, collateral, open and endarch vascular bundles constituted the primary vasculature of the stem (Fig. 6.4). A large centrally located pith was present.




Fig. 6.4. *Cycadeoidea*. Diagrammatic representation of T.S. stem showing primary structure.

The xylem and the phloem have been studied in detail by Wieland (1906) (Fig. 6.5 A, C) Most of the tracheids were rectangular in shape. They were scalariform. The tracheids of protoxylem were spiral. The secondary xylem and the secondary phloem were traversed by secondary medullary rays, which were either uniseriate or bi-seriate. Cambium was clearly visible.




Fig. 6.5. *Cycadeoidea wielandii*. A, T.S. stem passing near cambium; B, T.S. secondary wood; C, T.S. phloem showing thin-walled and thick-walled tracheids. (all after Wieland).

A leaf trace developed singly from the primary vascular strand. It divided into many mesarch strands upon entering into the cortex. At the place of its origin the leaf trace was C-shaped.

4. Reproductive Organs of *Cycadeoidea*:

The Bennettitalean reproductive organs are designated as “flowers “. The flower buds in the plants were present in the axil of leaf bases. As many on 500 flower buds were present on a single trunk in species such as *Cycadeoidea dartonii* (= *Monanthesia*

dartonii).

In several species of Cycadeoidea all the flower buds were present on a trunk at almost the same stage of development. Some palaeobotanists believe that such a plant might have flowered only once during its lifetime. Except a few species (e.g. *C. wielendii*) the flowers in Cycadeoidea were bisexual.

Hermaphrodite flower developed on a short pedicel. They were surrounded by as many as one hundred bracts, which were hairy and protective (Fig. 6.6). Flowers in different species were of different size. In *Cycadeoidea dartonii* they attained a length of about 2 cm and a diameter of about 1.5 cm while in *C. dacotensis* each flower was about 8 cm long and 3 cm in diameter.

In *C. dacotensis* the lower two-third portion of the floral axis had about 100-150 bracts. A whorl of stamens was present above the bracts. Each stamen was pinnately branched (Fig. 6.7) and each pinna had a double row of purse-shaped sporangia. Each sporangium resembled with a synangium. A conical floral axis was present just above the whorl of stamens. The entire compact structure resembled with a strobilus.




Fig. 6.6. *Cycadeoidea*. Diagrammatic representation of unexpanded strobilus. (after Wieland, 1906)




Fig. 6.7. *Cycadeoidea dacotensis*. Apical part with expanded and curved microsporophylls and a central conical axis. (after Wieland).

5. Microsporophyll in Cycadeoidea:

According to Wieland (1906, 1916), the androecium or pollen-bearing region consisted of about 20 pinnate, microsporophyll's. These were somewhat fixed or united at the base. Bean-shaped pollen capsules were arranged in two rows on each pinna of the

sporophyll.

These microsporophyll's remained folded round the gynoecium when young, but probably at maturity they expanded. Delevoryas (1963), however, opined that the microsporophyll's never expanded.

He further concludes that synangia-bearing structures, described as pinnae by Wieland (1906), were similar to the trabeculae. These trabeculae established a connection between outer and inner walls of the androecium. Pollen capsules or synangia were borne along these trabeculae.

Several (20-30) pollen sacs or microsporangia were present in a pollen capsule or synangium. The wall of a synangium consisted of outer palisade-like, thick-walled cells followed by thin-walled layer and then a tapetum.

The tapetum was not clearly demarcated (Figs. 6.8, 6.9). The pollen grains were oval in shape and measured up to 68 μ i in length. Multicellular pollen grains in Cycadeoidea




Fig. 6.8. *Cycadeoidea dacotensis*. L.S. sporangium showing stalk and two loculi with microspores. (after Wieland, 1906).

Fig. 6.9. *Cycadeoidea dacotensis*. T.S. of a synangium (after Wieland, 1906).

have been reported by Taylor (1973).

6. Gynoecium of Cycadeoidea:

The gynoecium receptacle was spherical or conical in shape. Hundreds of the stalked ovule along with an approximately equal number of inter-seminal scales were present

on the receptacle (Fig. 6.10). Each ovules was about 1 mm in length. The integument of the ovule was fused with the nucellus, except at the apex.

The ovule was orthotropous with a long micropylar beak. A pollen chamber and a nucellar beak was present in each ovule (Fig. 6.11) . The seed was somewhat elongated or oval in shape and possessed two cotyledons (Fig. 6.12) . Crepet and Delevoryas (1972) reported a linear tetrad in the nucellar region of Cycadeoidea.




Fig. 6.10. *Bennettites gibsonianus*. A female strobilus showing terminal seed with dicotyledonous embryos

Fig. 6.11. *Bennettites moneri*. L.S. seed. (after Wieland, 1906).




Fig. 6.12. *Bennettites albianus*. L.S. seed along with two surrounding interseminal scales. (after Stopes)

